


Wildlife Technical Report

Impact Evaluation with Wildlife Habitat Models and Evaluation of Cumulative Impacts to Wildlife in Southwestern Wyoming

Pinedale Anticline Oil and Gas Exploration and Development Project

Sublette County, Wyoming



Prepared By:

Bureau of Land Management
Pinedale Field Office
Pinedale, Wyoming

In Cooperation With:

U.S. Army Corps of Engineers
U.S. Forest Service
State of Wyoming

November, 1999

TABLE OF CONTENTS

SECTION	PAGE
PART I - WILDLIFE HABITAT MODELS	1
A. INTRODUCTION	1
B. BAYESIAN PROBABILITY MODELS	11
C. GIS ANALYSIS	12
Sources of Data	12
Hardware and Software Used in Analysis	12
Analytical Method	12
D. WILDLIFE MODELS	12
Species	12
Model Components	12
Modeling Cumulative Impacts	13
Pronghorn Winter Habitat Model	15
Mule Deer Winter Habitat Model	16
Sage Grouse Nesting Habitat Model	19
E. EXAMPLES OF MODEL APPLICATION IN CUMULATIVE IMPACT ANALYSIS	23
Pronghorn Winter Habitat Model	23
Mule Deer Winter Habitat Model	26
Sage Grouse Nesting Habitat Model	27
F. CUMULATIVE IMPACTS WITHIN THE PAPA	33
Potential and Existing Habitat Conditions in the PAPA	33
Model Simulations on the Pinedale Anticline Crest	33
Conclusion	36
PART II - CUMULATIVE IMPACTS TO WILDLIFE IN SOUTHWEST WYOMING	46
A. INTRODUCTION	46
B. CUMULATIVE IMPACT ANALYSIS AREA	46
Pronghorn	46
Mule Deer	46
Sage Grouse	46
C. LEVEL OF OIL AND GAS DEVELOPMENT	46
Pronghorn	46
Mule Deer	47
Sage Grouse	47
D. POSSIBLE EFFECTS ON POPULATION PARAMETERS - ANALYSIS	58
Big Game	58
Sage Grouse	59

TABLE OF CONTENTS
(continued)

SECTION	PAGE
E. POSSIBLE EFFECTS ON POPULATION PARAMETERS - RESULTS	61
Pronghorn	61
Mule Deer	63
Sage Grouse	66
LITERATURE CITED	67
APPENDIX A: Brief Outline Reviewing Probability with Application to Bayes' Theorem and Habitat Models	76
APPENDIX B: Estimation of Mule Deer and Pronghorn Maximum Survival Rates Between Ages 4-6 Months and 16-18 Months From Wyoming Game and Fish Department Herd Composition Data	81
APPENDIX C: Trends in Pronghorn Production Rates, Maximum Fawn Survival Rates and Total Winter Precipitation For Seven Antelope Herd Units in Southwest Wyoming	84
APPENDIX D: Trends in Mule Deer Production Rates, Maximum Fawn Survival Rates and Total Winter Precipitation For Six Mule Deer Herd Units in Southwest Wyoming	92

LIST OF TABLES

TABLE	PAGE
I.A-1. Summary of Wyoming Game and Fish Commission Mitigation Policy as it applies to wildlife species and wildlife habitats	7
I.A-2. Summary of primary and secondary impacts to wildlife expected on the PAPA due to any of the exploration/development scenarios	8
I.D-1. Model for pronghorn winter habitat	17
I.D-2. Model for mule deer winter habitat	20
I.D-3. Model for sage grouse nesting habitat	22
I.E-1. First example of applying the pronghorn winter habitat model to estimate potential habitat conditions, existing conditions, and cumulative conditions with a new project	24
I.E-2. Second example of applying the pronghorn winter habitat model to estimate potential habitat conditions, existing conditions, and cumulative conditions with a new project	25
I.E-3. First example of applying the mule deer winter habitat model to estimate potential habitat conditions, existing conditions, and cumulative conditions with a new project	28
I.E-4. Second example of applying the mule deer winter habitat model to estimate potential habitat conditions, existing conditions, and cumulative conditions with a new project	29
I.E-5. First example of applying the sage grouse nesting habitat model to estimate potential habitat conditions, existing conditions, and cumulative conditions with a new project	31
I.E-6. Second example of applying the sage grouse nesting habitat model to estimate potential habitat conditions, existing conditions, and cumulative conditions with a new project	32
I.F-1. Areas (acres and percent of total) of the PAPA predicted to be suitable habitats within various probability levels by three wildlife habitat models under potential and existing habitat conditions	34
I.F-2. Areas (acres and percent of total) of pronghorn winter ranges predicted to be suitable habitats within various probability levels in the PAPA under potential conditions, existing conditions, and different levels of well pad density, including 1 and 2 central production facilities (CPF), within four simulation areas of high exploratory interest on the Pinedale Anticline Crest	37
I.F-3. Areas (acres and percent of total) of mule deer winter ranges predicted to be suitable habitats within various probability levels in the PAPA under potential conditions, existing conditions, and different levels of well pad density, including 1 and 2 central production facilities (CPF), within four simulation areas of high exploratory interest on the Pinedale Anticline Crest	38
I.F-4. Areas (acres and percent of total) of sage grouse nesting areas predicted to be suitable habitats within various probability levels in the PAPA under potential conditions, existing conditions, and different levels of well pad density, including 1 and 2 central production facilities (CPF), within four simulation areas of high exploratory interest on the Pinedale Anticline Crest	39
II.A-1. Summary of oil and gas development projects previously or currently being analyzed under NEPA in southwest Wyoming as of December, 1998	48
II.C-1. Oil and gas developments in southwest Wyoming since 1984 that coincide with seasonal ranges used by pronghorn in 7 Antelope Herd Units	53

LIST OF TABLES
(continued)

TABLE	PAGE
II.C-2. Oil and gas developments in southwest Wyoming since 1984 that coincide with seasonal ranges used by mule deer in 7 Mule Deer Herd Units	55
II.C-3. Oil and gas developments in southwest Wyoming since 1984 that coincide with sage grouse lek locations in 2 Upland and Small Game Management Areas (USGMA)	57
II.E-1. Pronghorn population parameters in southwest Wyoming herd units before and 1 year after major oil/gas project NEPA decision documents were issued by BLM for projects coinciding with each herd unit. Fawn production rate (number of fawns divided by number of does 2 years or older counted pre-harvest, before the hunting season) and maximum fawn survival rate (estimated for the period from age 4 months to age 16 months) were averaged for the time periods shown (with standard errors, SE)	62
II.E-2. Comparisons of total winter precipitation, pronghorn fawn production rates and survival rates in 7 Antelope Herd Units during the severe winter of 1983-84 and winters of the 1990's with lowest fawn production and/or survival rate estimates	62
II.E-3. Mule deer population parameters in southwest Wyoming herd units before and 1 year after major oil/gas project NEPA decision documents were issued by BLM for projects coinciding with each herd unit. Fawn production rate (number of fawns divided by number of does 2 years or older counted post-harvest, after the hunting season) and maximum fawn survival rate (estimated for the period from age 6 months to age 18 months) were averaged for the time periods shown (with standard errors, SE)	64
II.E-4. Comparisons of total winter precipitation, mule deer fawn production rates and survival rates in 6 Mule Deer Herd Units (no data was collected in the Chain Lakes Herd Unit) during the severe winter of 1983-84 and winters of the 1990's with lowest fawn production and survival rate estimates	64
II.E-5. Numbers of active, inactive and unknown activity status sage grouse leks that have at least 1 oil or gas well within the specified distances in 2 Upland and Small Game Management Areas (USGMA)	66

LIST OF FIGURES

FIGURE	PAGE
I.A-1. General location of the Pinedale Anticline Project Area	3
I.A-2. Pronghorn seasonal ranges in the Project Area	4
I.A-3. Mule deer seasonal ranges in the Project Area	5
I.A-4. Sage grouse leks in the Project Area	6
I.D-1. Areas of high geologic interest used to simulate the effects of exploration/development on the Pinedale Anticline Crest	14
I.F-1. Pronghorn winter habitat within the PAPA with four probability categories for being suitable. The PAPA was evaluated under potential conditions	40
I.F-2. Pronghorn winter habitat within the PAPA with four probability categories for being suitable. The PAPA was evaluated under existing conditions	41
I.F-3. Mule deer winter habitat within the PAPA with four probability categories for being suitable. The PAPA was evaluated under potential conditions	42
I.F-4. Mule deer winter habitat within the PAPA with four probability categories for being suitable. The PAPA was evaluated under existing conditions	43
I.F-5. Sage grouse nesting habitat within the PAPA with four probability categories for being suitable. The PAPA was evaluated under potential conditions	44
I.F-6. Sage grouse nesting habitat within the PAPA with four probability categories for being suitable. The PAPA was evaluated under existing conditions	45
II.A-1. Existing and potential oil and gas projects in southwest Wyoming	49
II.B-1. Antelope Cumulative Impact Analysis Area	50
II.B-2. Mule Deer Cumulative Impact Analysis Area	51
II.B-3. Sage Grouse Cumulative Impact Analysis Area	52
II.D-1. Relationship of overwinter fawn and adult mule deer mortality rates as nonlinear functions of total winter precipitation, November through March, since 1992 in the Sublette Mule Deer Herd Unit. The relationship of winter precipitation to fawn mortality rate is significant ($r^2 = 0.84$, $p = 0.004$) as is the relationship to adult mortality rate ($r^2 = 0.61$, $p = 0.038$). Brackets are 90% confidence intervals on each estimate. Winter precipitation data were averaged for NOAA cooperators at Pinedale, Daniel Fish Hatchery, and Boulder Rearing Station	60
II.E-1. Maximum fawn survival rates in the Sublette Mule Deer Herd Unit decrease slightly with increasing winter precipitation from 1979 to 1994 (solid line, $y = 0.599 - 0.072 x$, $p = 0.049$). Maximum fawn survival rates from 1995 to 1998 decrease sharply with increasing winter precipitation (dashed line, $y = 1.048 - 0.228 x$, $p = 0.068$)	65